

---

## Thesis subject

Name of the laboratory: LAM/IRAM

Thesis advisor: **Alexandre Beelen**

Email and address: [alexandre.beelen@lam.fr](mailto:alexandre.beelen@lam.fr)

Tel: 04 95 04 41 04

Co-advisor: Guilaine Lagache / [guilaine.lagache@lam.fr](mailto:guilaine.lagache@lam.fr) / 04 95 04 41 77

Subject's title: Mapping the dusty star formation at high redshift with the NIKA2 Cosmological Legacy Survey

Subject description:

Among the potentially most important results of cosmology over the last 15 years has been the discovery of the importance of dust obscured star formation. While dusty galaxies are rare in the local Universe ( $\sim 1\%$ ), they are 10 to 100 times more numerous at higher redshift ( $z > 1$ ). For these galaxies, a large fraction of the light produced by stars reaches us in the range 100-2000 microns, after being reprocessed by dust. These galaxies become so important that they dominate the census of star formation in the Universe at  $1 < z < 3.5$ . At higher redshift ( $z > 4$ ), quantifying the star formation rate density is a challenging endeavour. Today, most of the high- $z$  measurements are based on UV emission emerging from the galaxies. However, these UV observations trace the un-obscured star formation and do not allow to directly measure the star formation from dusty galaxies. One of the crucial questions is therefore to quantify the role of these galaxies in the overall star formation budget at  $z > 3.5$ .

To reach this goal, new deep and unbiased millimetre surveys are mandatory. Such surveys must combine high sensitivity and wide area, with good angular resolution to reduce confusion noise. Confusion is the ultimate limit of cosmological survey; it is inherent of deep surveys on a single dish telescope in the millimetre and sub-millimetre domain. Today, only a new generation of millimetre camera on a large telescope, with a high mapping speed, can fulfil the conditions to fully map the obscured star formation at high- $z$ . This is one of the objectives of the cosmological surveys made with the NIKA2 camera at 1.2 and 2mm. Built at the Institut Néel in Grenoble, NIKA2 is installed on the 30m IRAM telescope and is now open to the community. At LAM, our group is responsible of a Guaranteed Time Large Program, the "NIKA2 Cosmological Legacy Survey", with 300 hours of observations spread over 4 years on the GOODS North and COSMOS fields. The observations already started in October 2017.

The thesis objectives are a) to obtain a complete view of the star formation rate at high- $z$ ; b) to bring new constraints on the evolution of galaxies (by probing for example the star formation in "normal" galaxies at high- $z$  and determining if these galaxies follow the main

sequence of star formation seen in the Universe at  $z < 2$  or if the starburst activity induced by galaxies interaction and fusion play an important role); c) to measure the distribution of star formation in large-scale structures with clustering measurements.

The successful PhD candidate will participate to the observation campaigns on the IRAM 30m at Pico Veleta (Spain), and will also contribute to the data processing and analysis. She/he will extract the sources using multi-wavelength source extraction codes. Novel extraction techniques could also be explored, such as image segmentation based on Deep Learning algorithm. Using counterparts of detected millimetre sources using ancillary data, she/he will analyse their spectral energy distribution to measure stellar masses and star formation rate. Once the observations and galaxy detections are fully characterized, and translated in terms of source counts, she/he will directly use them to constrain the history of dusty star formation by fitting for example our evolutionary model of galaxy evolution.

She/he will also help us to define the best strategy for follow up observations, in particular with IRAM NOEMA and/or ALMA, to measure the redshift and gas mass of the most representative galaxies. Such measurements will be used to relate the star formation efficiency to other galaxy properties (mass, merger state, kinematics), to learn what regulates their star formation. Finally, she/he could be involved in the study of the clustering of these objects, that will put particularly strong constraints on models for the evolution of large scale structure, giving the link between star formation and dark matter halos mass.

This thesis, co-funded by IRAM, offers the unique opportunity to participate to a pioneering observational project, within a dedicated team at LAM. Members of the NIKA2 deep field surveys include M. Béthermin, V. Buat and D. Burgarella, in addition to the PIs G. Lagache & A. Beelen, at LAM as well as many others over the globe.

#### Bibliography:

##### NIKA2:

<http://lpsc.in2pf.fr/NIKA2Sky> & <http://ipag.osug.fr/nika2>

Adam R., et al., 2018, A&A, 609, A115

##### (Sub-)millimeter deep field observations:

Lindner R.R., et al., 2011, ApJ, 737, 83

Geach J.E., et al., 2017, MNRAS, 465, 1789

Dunlop J.S., et al., 2017, MNRAS, 466, 861

##### Dusty galaxies Reviews:

Lagache G. , et al. 2005 AARA&A 43, 727

Casey C.M., Narayanan D., Cooray A., 2014, PhR, 541, 45